









## INJECTION INSTRUCTIONS

1. Wash hands.
2. Take out medication and check name and expiration date.
3. Draw up specified amount of medication. (Refer to specific medication instructions for mixing directions.)
4. Remove air from syringe. The easiest way to do this is to hold the syringe upright and flick it. The air will form a bubble closest to the needle and can then be pushed out of the syringe.
5. To locate injection site, see illustration below.

<b>SUB Q INJECTIONS</b>		
Either side of fatty area of umbilicus 	Fatty area of upper outer arm 	Fatty area of upper thigh 
<b>IM INJECTIONS</b>		
Upper outer quadrant of buttocks 	Upper outer thigh 	Deltoid area upper arm 

6. Cleanse site with alcohol.
7. **For IM:** With one hand pull the skin tight. Holding the syringe like a pencil in the other hand, use a wrist like action to quickly insert needle into muscle at a 90° angle (all the way to plastic hub). Holding needle in place, aspirate (pull back plunger) to check for blood. If you see blood, remove needle and start over with clean needle.  
**For SubQ:** With one hand, pinch the skin between the thumb and fore-finger. Holding the syringe like a pencil with the other hand, insert the small needle just under the skin at a 45° angle.
8. Inject medication.
9. Remove needle and massage area.
10. Do not re-cap needle and dispose into red puncture-proof bio-hazard container.
11. Alternate sites of injections.